

University of Wollongong Research Online

University of Wollongong Thesis Collection
1954-2016

University of Wollongong Thesis Collections

2008

A morphodynamic model of atoll-islet evolution

Stephen J. Barry
University of Wollongong

Follow this and additional works at: <https://ro.uow.edu.au/theses>

University of Wollongong

Copyright Warning

You may print or download ONE copy of this document for the purpose of your own research or study. The University does not authorise you to copy, communicate or otherwise make available electronically to any other person any copyright material contained on this site.

You are reminded of the following: This work is copyright. Apart from any use permitted under the Copyright Act 1968, no part of this work may be reproduced by any process, nor may any other exclusive right be exercised, without the permission of the author. Copyright owners are entitled to take legal action against persons who infringe their copyright. A reproduction of material that is protected by copyright may be a copyright infringement. A court may impose penalties and award damages in relation to offences and infringements relating to copyright material.

Higher penalties may apply, and higher damages may be awarded, for offences and infringements involving the conversion of material into digital or electronic form.

Unless otherwise indicated, the views expressed in this thesis are those of the author and do not necessarily represent the views of the University of Wollongong.

Recommended Citation

Barry, Stephen J, A morphodynamic model of atoll-islet evolution, PhD thesis, School of Earth and Environmental Sciences, University of Wollongong, 2008. <http://ro.uow.edu.au/theses/378>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

NOTE

This online version of the thesis may have different page formatting and pagination from the paper copy held in the University of Wollongong Library.

UNIVERSITY OF WOLLONGONG

COPYRIGHT WARNING

You may print or download ONE copy of this document for the purpose of your own research or study. The University does not authorise you to copy, communicate or otherwise make available electronically to any other person any copyright material contained on this site. You are reminded of the following:

Copyright owners are entitled to take legal action against persons who infringe their copyright. A reproduction of material that is protected by copyright may be a copyright infringement. A court may impose penalties and award damages in relation to offences and infringements relating to copyright material. Higher penalties may apply, and higher damages may be awarded, for offences and infringements involving the conversion of material into digital or electronic form.

A Morphodynamic Model of Atoll-islet Evolution

A thesis submitted in fulfilment of the requirements for the award of the
degree

Doctor of Philosophy

from

The University of Wollongong

by

Stephen J. Barry BSc (Hons)

School of Earth and Environmental Sciences

Faculty of Science

2008

Certification

I, Stephen J. Barry, declare that this thesis, submitted in fulfillment of the requirements for the award of Doctor of Philosophy, in the School of Earth and Environmental Sciences, University of Wollongong, is wholly my own work unless otherwise referenced or acknowledged. The document has not been submitted for qualifications at any other academic institution.

Stephen J. Barry

Date:

"There are two kinds of truths: those of reasoning and those of fact"

Gottfried Wilhelm Leibniz

"All models are wrong, but some are useful"

George E. P. Box

Acknowledgements

This work would not exist at all without the contributions and support of many friends, colleagues and family. I owe many thanks to Dr Peter Cowell, whose knowledge, guidance, and enthusiasm, has been fundamental to the progression of this research and to my education in both academic and coastal behavior.

Thanks are also due to my supervisor Colin Woodroffe for providing the initial funding and motivation for this research. I am indebted to the staff of the School of Earth and Environmental Sciences for their support during this period. Thanks also to Paul Carr, Laurie Chisholm, and Marji Puotinen for providing advice and perspective; John Reid, Mike Robinson, Heidi Brown, and John Marthick for technical support. I am pleased to acknowledge and thank Dr Peter Eklund, Dr Mark Sifer, and Dr Reza Zamani from the Faculty of Commerce for providing advice, employment, and encouragement during my time at the University of Wollongong.

Support from fellow students, past and present, has been essential during this project; thanks to Paul Grevenitz, Ava Simms, Paolo Aballe, Gareth Davies, Kevin Pucillo, and George Susino. Thanks also to Peredur Jones who was a reliable companion for extramural amusements.

Most of all I would like to thank my family for their support throughout these years of study. I am especially pleased to be able to thank my wife, Lynne; without her unwavering perseverance and commitment this thesis would not have been completed.

Table of Contents

Certification.....	ii
Acknowledgements	iv
Table of Contents	v
List of Figures	viii
List of Tables.....	x
List of Notation	xi
List of Abbreviations.....	xiv
List of Publications	xv
Abstract	xvi
1 Introduction	1
1.1 General Introduction	2
1.2 Atoll-islet Morphology	3
1.3 Modeling Coastal Behavior.....	5
1.4 The Sediment Allocation Model	8
1.5 Thesis Organization	9
2 A Morphodynamic Model of Atoll-Islet Evolution	11
2.1 Abstract	12
2.2 Introduction	13
2.2.1 Purpose of the Model	16
2.3 The Sediment Allocation Model	17
2.4 Model Parameters.....	22
2.4.1 A Hypothetical Atoll-islet.....	24
2.5 Parameter Estimation Using Inverse Techniques	30
2.5.1 Optimization Constraints	32
2.5.2 Parameter Estimation	34
2.6 Results	37
2.7 Discussion	39

2.8	Conclusion	41
3	Growth-Limiting Size of Atoll-Islets: Morphodynamics in Nature.....	43
3.1	Abstract	44
3.2	Introduction	45
3.3	Methods.....	49
3.3.1	The Sediment Allocation Model	50
3.3.2	Field Data	53
3.4	Boundary Conditions and Cell Definition for Transect Data-model	55
3.4.1	Choice of Isochron Pattern for Transect Data-model	57
3.4.1.1	Progradational data-model	58
3.4.1.2	Hybrid Data-model.....	59
3.5	Results	63
3.5.1	Data-model Comparison	63
3.5.2	Parameter Estimation	65
3.5.3	A Simulation of Islet Transect Development.....	67
3.6	Discussion	71
3.6.1	Simulating Islet Growth	74
3.6.2	Sampling Strategy	76
3.7	Conclusion	78
4	Atoll-Islet Morphology and Energy Exposure: Comparative Metrics.....	80
4.1	Abstract	81
4.2	Introduction	82
4.2.1	Models of Atoll Development.....	84
4.2.2	Rationale for a Comparative Metric.....	90
4.2.3	Measuring Islet Location.....	91
4.3	Materials and Methods	94
4.3.1	Field Setting	94
4.3.2	Energy Exposure	94
4.3.2.1	Exposure Regions	98
4.3.2.2	Antecedent Morphology.....	99
4.3.2.3	Measurement of Islet Location.....	102
4.4	Results	104

4.5	Discussion	107
4.5.1	Energy Exposure and Islet Location	107
4.5.2	Cocos_E and Marakei_W	107
4.5.3	Marakei_W and Tarawa_SW	110
4.5.4	Tarawa_N and Marakei_E	111
4.6	Conclusion	112
4.6.1	Islet Response to SLR	114
4.7	Acknowledgements	116
5	Conclusions	117
5.1	Introduction	118
5.2	Directions for Future Research	119
5.3	Conclusion	121
6	References	123
	Appendix A	140
	Appendix B	141
	Appendix C	142

List of Figures

Figure 1.1 Idealized representation of an atoll-islet environment.....	4
Figure 1.2 Response of sediment sequestration to atoll-islet evolution.....	7
Figure 2.1 The Sediment Allocation Model.....	18
Figure 2.2 The forward model of islet development.....	19
Figure 2.3 Evolution of the islet sediment sequestration ratio.....	23
Figure 2.4 An example of the physical setting of an atoll islet.....	25
Figure 2.5 The sediment allocation function generated using the parameter values $b = 200$, $r = 10$, and $V_{\max} = 600,000$	26
Figure 2.6 Output from the forward model shows the growth of an atoll isle.....	26
Figure 2.7 The sediment allocation function when the r parameter is altered.....	27
Figure 2.8 The effect on the SAM for different values for the parameter r	28
Figure 2.9 The sediment allocation function when the b parameter is altered	29
Figure 2.10 The effect on the SAM for different values for the parameter b	30
Figure 2.11 Graphs of the averaged estimated parameters	38
Figure 2.12 The output produced by the forward model	39
Figure 3.1 Schematic scenarios of island formation	47
Figure 3.2 A conceptual model of aggradational reef-island growth.....	48
Figure 3.3 An example of a sediment allocation function	52
Figure 3.4 Location of the field sites	54
Figure 3.5 Transect data for Cocos (Keeling) Island	54
Figure 3.6 Transect data for Makin Island	55
Figure 3.7 A conceptual model of the processes contributing to reef-island.....	56
Figure 3.8 The modified transect images.....	60

Figure 3.9 The elliptical isochrons used to calculate the transect volumes	62
Figure 3.10 Graphs of transect development based on evolution by lateral accretion evolution by hybrid accretion.....	64
Figure 3.11 Sediment allocation functions for the study sites	66
Figure 3.12 Output from the SAM using estimated parameters	66
Figure 3.13 Simulation of reef-island transect development	70
Figure 3.14 An interpretation of the effect that the SAM parameters	72
Figure 4.1 Representation of an atoll-islet as morphological sub-units.....	88
Figure 4.3 The location of the sites showing the regions of the islets sampled.....	95
Figure 4.4 The Tarawa_N region contains significant conglomerate deposits.....	100
Figure 4.5 The sampling method used to measure the distance from the reef rim to the atoll islet.....	103
Figure 4.6 Boxplot of reef-rim to islet-beach distance	106

List of Tables

Table 2.1 Parameters used to represent an atoll-islet environment.....	25
Table 2.2 Constraint values for the inversion algorithm.....	32
Table 2.3 Selected Sink 1 volumes from the forward model.....	35
Table 2.4 The average values of estimated parameters.....	37
Table 3.1 Estimated parameters for the SAM.....	65
Table 3.2 Measurements used to model the reef-island profile.	68
Table 4.1 Energy exposure variables considered for the four study sites.....	97
Table 4.2 Ranking of off-reef energy-exposure by region.....	99
Table 4.3 Summary measurements of islet location for sampled islet regions.....	105

List of Notation

a	height at the transect origin.
b	sediment transport impedance term. The linear parameter in the logistic function representing transport gradients relevant to sediment sequestration by the island.
c	$\ln(h/a)L$.
D	a <i>Sloss</i> variable; the dispersal factor.
i	increment counter.
$1-f(v/V_{\max})$	island sediment sequestration ratio. The proportion of available sediment sequestered by the reef-island.
$f(v/V_{\max})$	the proportion of available sediment that bypasses the reef-island.
$f1_error$	$1-f(v/V_{\max})$ when $v/V_{\max} = 1$. When the accommodation volume is filled no sediment is sequestered by the island. This means that, ideally, the value of $1-f(1) = 0$, however, since $0 < f(v/V_{\max}) < 1$, this will never be the case. Instead, the $f1_error$ can be used as an error term in the nonlinear optimization routine.
h	ridge height.
L	length measurement.
m	meter.
p	the perturbation amount calculated as $p = \tilde{v}(i) \times p_{\max} \times x$. This value was used to perturb the exact value, $\tilde{v}(i)$, during testing of the inversion procedure. For example, if $\tilde{v}(i) = 100$, $p_{\max} = 0.1$, and $x = -0.5$, then $p = 100 \times 0.1 \times -0.5 = -5$, then the value of $\tilde{v}(i)$ used in the parameter recovery process is $\tilde{v}(i) = \tilde{v}(i) - 5 = 100 + 5 = 105$.

p_{\max}	the maximum perturbation ratio. If the measured value was 100 and the maximum perturbation ratio was 0.1, then, the maximum absolute value of the perturbation is $100 \times 0.1 = 10$.
Q	a <i>Sloss</i> variable; the quantity of sediment available for deposition.
r	transport-impedance sensitivity to morphology. The nonlinear parameter in the logistic function that represents morphodynamic feedback parameter, i.e. the response of sediment transport processes to island growth.
R	a <i>Sloss</i> variable; the accommodation available for sediment deposition.
s	rate of sediment production; seconds.
s_i	sediment produced during step i of the forward model.
t	time.
t_0	$t = 0$.
\tilde{v}	vector of island volume measurements.
v_0	initial volume of the island at $t = 0$. In the solution to the logistic equation $b = (1/t_0) - 1$, therefore $t_0 \neq 0$, and $v_0 \neq 0$. This implies that the island forms as the result of a pre-existing perturbation of the sediment transport field. For the purposes of testing it was assumed that the initial island volume was 0 m^3 .
\tilde{v}'	vector of island volumes calculated using the forward model.
V_{\max}	accommodation volume, the maximum volume available for island development.
$w()$	weighting function. This is used to magnify the <i>fl_error</i> value so that it is significant when compared to the calculated error term in the optimization process.

x	unit of length; a random number in the range $[-1, 1]$.
y	unit of length.
yr	year.

List of Abbreviations

ECD	Effective conglomerate distance
ENSO	El Niño Southern Oscillation
GIS	Geographic Information System
IQR	Interquartile range
LD	Lagoonside deposition
OD	Oceanside deposition
SAM	Sediment Allocation Model
SLR	Sea-level rise
VBA	Visual BASIC for Applications

List of Publications

- Barry, S.J.**, Cowell, P.J. and Woodroffe, C.D., 2007. A morphodynamic model of reef-islet development on atolls. *Sedimentary Geology*, 197(1-2): 47-63.
- Barry, S.J.**, Cowell, P.J. and Woodroffe, C.D., 2008. Growth-limiting size of atoll-islets: Morphodynamics in nature. *Marine Geology*, 247(3-4): 159-177.
- Barry, S.J.**, Cowell, P.J. and Woodroffe, C.D., Atoll-islet morphology and energy exposure: comparative metrics, *Manuscript submitted*.

Abstract

A morphodynamic model of atoll-islet evolution, the Sediment Allocation Model (SAM), was developed based on the assumption that islets are equilibrium landforms. The assumption that islets are equilibrium landforms implies that the volume of sediment sequestered in an islet reaches stability when considered over the time-scale of islet evolution. Stability of the islet implies that the maximum volume of an islet (accommodation) can be quantified. Morphodynamic feedback between the islet and sediment transport processes is manifest in the rate of sediment sequestration by the islet. Initial increase of islet volume is rapid, sediment sequestration slows as islet volume increases, and stops when accommodation is full.

The rate of islet development is constrained by antecedent morphology, sediment supply, and sediment transport. The constraints and accommodation for islet evolution are difficult to measure directly. Estimates of the accommodation, sediment supply, the effect of antecedent morphology, and the morphodynamic feedback between islet volume and sediment sequestration are generated using measurements of islet volume in an inversion algorithm. Experimentation carried out using a hypothetical test-case indicated that errors in the measurements of islet volume used in the inversion algorithm would result in a damping of the morphodynamic feedback and overestimation of both sediment supply and accommodation.

The SAM was implemented using published radiocarbon-dated samples for surveyed islets transects. A data-model template was developed to incorporate the published data into the SAM. In the data-model template an atoll-islet was represented by a

combination of morphological sub-units with sub-unit sequestration dominated by either lagoon processes or ocean processes. The input to the SAM was a series of sediment volumes representing islet evolution. Sediment volumes were calculated using isochron patterns that represent a hybrid of lateral accretion and vertical accretion, recognized modes of sediment accumulation on atoll-islets. Sediment volume measurements calculated using the hybrid-accretion patterns were consistent with the assumption of morphodynamic feedback between islet morphology and sediment accretion processes.

Comparative metrics are required to test and refine models of islet evolution, however, suitable comparative data are not currently available. The hybrid-accretion pattern of islet evolution formed the basis for the development of a metric to determine islet location on the reef platform. Islet location was measured on four atolls as the distance from the reef crest to the oceanward beach-toe. The results of the measurement procedure were evaluated in the context of three factors that influence islet evolution: energy-exposure, sediment supply, and antecedent morphology. The results indicate that the distance from the reef crest to the oceanward beach-toe is a measurable response to the influence of the three factors.